

Evaluate Different Trig Expressions Worksheet Answer Key PDF

Evaluate Different Trig Expressions Worksheet Answer Key PDF

Disclaimer: The evaluate different trig expressions worksheet answer key pdf was generated with the help of StudyBlaze AI. Please be aware that AI can make mistakes. Please consult your teacher if you're unsure about your solution or think there might have been a mistake. Or reach out directly to the StudyBlaze team at max@studyblaze.io.

Part 1: Building a Foundation

Which of the following is the reciprocal of the sine function?

undefined. A) Cosine

undefined. A) Secant

undefined. A) Cosecant ✓

undefined. A) Tangent

The reciprocal of the sine function is cosecant.

Select all the correct Pythagorean identities.

undefined. A) $\sin^2(\theta) + \cos^2(\theta) = 1$ ✓

undefined. A) $1 + \tan^2(\theta) = \sec^2(\theta)$ ✓

undefined. A) $\sin(\theta)\cos(\theta) = 1$

undefined. A) $1 + \cot^2(\theta) = \csc^2(\theta)$ ✓

The correct Pythagorean identities are $\sin^2(\theta) + \cos^2(\theta) = 1$, $1 + \tan^2(\theta) = \sec^2(\theta)$, and $1 + \cot^2(\theta) = \csc^2(\theta)$.

Define the tangent function in terms of sine and cosine.

The tangent function is defined as $\tan(\theta) = \sin(\theta)/\cos(\theta)$.

List the trigonometric values for $\sin(30^\circ)$, $\cos(45^\circ)$, and $\tan(60^\circ)$.

1. $\sin(30^\circ)$

1/2

2. $\cos(45^\circ)$

$\sqrt{2}/2$

3. $\tan(60^\circ)$

$\sqrt{3}$

The values are $\sin(30^\circ) = 1/2$, $\cos(45^\circ) = \sqrt{2}/2$, and $\tan(60^\circ) = \sqrt{3}$.

Part 2: comprehension and Application

Which identity can be used to simplify $\sin(\alpha + \beta)$?

undefined. A) $\sin(\alpha)\cos(\beta) + \cos(\alpha)\sin(\beta)$ ✓

undefined. A) $\cos(\alpha)\cos(\beta) - \sin(\alpha)\sin(\beta)$

undefined. A) $\tan(\alpha) + \tan(\beta)$

undefined. A) $\sin^2(\alpha) + \cos^2(\beta)$

The identity used to simplify $\sin(\alpha + \beta)$ is $\sin(\alpha)\cos(\beta) + \cos(\alpha)\sin(\beta)$.

Which of the following are correct values for special angles?

undefined. A) $\sin(45^\circ) = \sqrt{2}/2$ ✓

undefined. A) $\cos(60^\circ) = 1/2$ ✓

undefined. A) $\tan(90^\circ) = 1$

undefined. A) $\sin(90^\circ) = 1$ ✓

The correct values for special angles are $\sin(45^\circ) = \sqrt{2}/2$, $\cos(60^\circ) = 1/2$, and $\sin(90^\circ) = 1$.

Explain the significance of the unit circle in trigonometry.

The unit circle provides a geometric representation of trigonometric functions, allowing for the visualization of angles and their corresponding sine and cosine values.

Solve the equation $2\sin(x) - 1 = 0$ for x in the interval $[0^\circ, 360^\circ]$.

The solutions are $x = 30^\circ$ and $x = 150^\circ$.

Part 3: Analysis, Evaluation, and Creation

Which graph represents a function with a period of π ?

undefined. A) $y = \sin(x)$

undefined. A) $y = \cos(x)$

undefined. A) $y = \tan(x)$ ✓

undefined. A) $y = \sec(x)$

The graph that represents a function with a period of π is $y = \tan(x)$.

Analyze the following expressions and select those that are equivalent to $\tan(\theta)$.

undefined. A) $\sin(\theta)/\cos(\theta)$ ✓

undefined. A) $1/\cot(\theta)$ ✓

undefined. A) $\cos(\theta)/\sin(\theta)$

undefined. A) $\sec(\theta)\csc(\theta)$

The expressions equivalent to $\tan(\theta)$ are $\sin(\theta)/\cos(\theta)$ and $1/\cot(\theta)$.

Compare and contrast the graphs of $y = \sin(x)$ and $y = \cos(x)$, focusing on amplitude and phase shift.

Both graphs have the same amplitude of 1, but the sine graph has a phase shift of $\pi/2$ compared to the cosine graph.

Which of the following scenarios best describes an application of trigonometry in real life?

undefined. A) Calculating the area of a rectangle

undefined. A) Determining the height of a building using its shadow ✓

undefined. A) Measuring the volume of a cylinder

undefined. A) Counting the number of sides in a polygon

Determining the height of a building using its shadow is a common application of trigonometry.

Evaluate the following statements and select those that correctly describe the properties of inverse trigonometric functions.

undefined. A) $\arcsin(x)$ is defined for all real numbers.

undefined. A) $\arccos(x)$ has a range of $[0, \pi]$. ✓

undefined. A) $\arctan(x)$ is defined for all real numbers. ✓

undefined. A) $\arcsin(x)$ has a range of $[-\pi/2, \pi/2]$. ✓

The correct statements are $\arccos(x)$ has a range of $[0, \pi]$, $\arctan(x)$ is defined for all real numbers, and $\arcsin(x)$ has a range of $[-\pi/2, \pi/2]$.

Design a real-world problem that involves using trigonometric identities to find an unknown angle or side in a triangle. Describe the problem and outline the steps to solve it.

An example problem could involve finding the height of a tree using the angle of elevation and distance from the tree.